

## Anti-LSD R12301 SPTN-5

### Product overview

---

<b>Catalog number</b>	140027
<b>Specificity</b>	Antibody recognizes lysergic acid diethylamide (LSD)
<b>Description</b>	Recombinant mouse IgG <sub>1</sub> antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	18 months from manufacturing at 2–8 °C
<b>Subclass</b>	IgG <sub>1</sub>
<b>Analyte description</b>	Lysergic acid diethylamide (LSD) is a semi-synthetic hallucinogen.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/- 10%) (A280 nm, 1 mg/ml, 1 cm = 1.6)
<b>Immunoreactivity</b>	80–120% compared to the reference sample in an FIA test
<b>IEF Profile</b>	6.3–6.9
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	$1.6 \times 10^6$ 1/Ms
<b>Dissociation rate constant</b>	$1.2 \times 10^{-4}$ 1/s
<b>Affinity constant</b>	$K_A = 1.4 \times 10^{10}$ 1/M; $K_D = 9.5 \times 10^{-11}$ M (= 0.10 nM)
<b>Determination method</b>	BLI (Octet RED96e)
<b>Determination antigen</b>	Lysergic acid diethylamide (LSD)-BSA, EastCoast Bio LA783



#### Legal disclaimer

<b>Cross-reactivities</b>	Recognizes 2-Oxo-3-hydroxy-LSD													
<b>Epitope</b>	N/D													
<b>Pair recommendations</b>	Lysergic acid diethylamide (LSD)-BSA Antigen LA783													
	Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.													
<b>Platforms tested</b>	FIA													
<b>Antigens tested</b>	Please see pair recommendations above.													
<b>Product stability</b>	<table><thead><tr><th>TEMPERATURE, TIME</th><th>RESULT</th></tr></thead><tbody><tr><td>-70 °C, 21 days</td><td>OK</td></tr><tr><td>-20 °C, 21 days</td><td>OK</td></tr><tr><td>+4 °C, 21 days</td><td>OK</td></tr><tr><td>+35 °C, 21 days</td><td>OK</td></tr><tr><td>+45 °C, 7 days</td><td>OK</td></tr></tbody></table>	TEMPERATURE, TIME	RESULT	-70 °C, 21 days	OK	-20 °C, 21 days	OK	+4 °C, 21 days	OK	+35 °C, 21 days	OK	+45 °C, 7 days	OK	
TEMPERATURE, TIME	RESULT													
-70 °C, 21 days	OK													
-20 °C, 21 days	OK													
+4 °C, 21 days	OK													
+35 °C, 21 days	OK													
+45 °C, 7 days	OK													
	Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.													
<b>Miscellaneous</b>	-													
<b>References</b>	-													



#### Legal disclaimer